

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 October 2002 (17.10.2002)

PCT

(10) International Publication Number
WO 02/081614 A1

(51) International Patent Classification⁷: **C11D 3/50**,
A61K 7/32, C11D 3/00, A61K 7/46

(21) International Application Number: PCT/GB02/01609

(22) International Filing Date: 4 April 2002 (04.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0108657.8 6 April 2001 (06.04.2001) GB

(71) Applicant (*for all designated States except US*): **QUEST INTERNATIONAL B.V.** [NL/NL]; Huizerstraatweg 28, NL-1411 GP Naarden (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **PERRING, Keith**, Douglas [GB/GB]; 14 Malvern Road, Ashford, Kent TN24 8HS (GB). **GOODACRE, Terence** [GB/GB]; Southlands, Ashford Road, High Halden, Kent TN26 3BP (GB).

(74) Agent: **KEITH W NASH & CO**; 90-92 Regent Street, Cambridge CB2 1DP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) *for US only*

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PERFUME COMPOSITIONS**

(57) Abstract: This invention is concerned with perfume compositions comprising: a) nitrile perfumery material of general formula RCN where R is a hydrocarbyl radical of formula C_xH_y, x is an integer in the range 9 to 12, and y is an integer in the range 15 to 25; and b) hydrocarbon perfumery material having a boiling point of less than about 185 °C and an octanol-water partition coefficient of at least 4.0 (in logarithmic form), wherein a) and b) together comprise at least 20 % by weight of the total weight of the composition. Perfume compositions in accordance with the present invention provide a good deodorant effect several hours after application to a surface such as fabric or skin.



WO 02/081614 A1

Title: Perfume CompositionsField of the Invention

This invention relates to perfume compositions and to products containing such perfume compositions. The invention also concerns the use of perfume compositions to provide a deodorant effect.

Background to the Invention

A highly desired attribute of many consumer products is that they include fragrances which demonstrate long-lasting effect e.g. following application to a surface. Several approaches have been used to deliver this property from a fragrance. These include, for example, the inclusion of a high boiling, low odour, fixative material to extend fragrance lifetime as described in US 6172037, or the encapsulation of fragrance inside capsules which rupture or degrade following application to a surface as described in EP 0397246. Both of these approaches, however, may add to the cost of the fragrance and/or may result in a composition which is unsuitable for use in a variety of consumer products, possibly requiring further developmental work.

One alternative approach, as described in WO97/30689 and US 5500154, lies in the formulation of perfumes that provide a long lasting aesthetic benefit with a minimum amount of material. These perfumes have been described as 'enduring perfumes'. Enduring perfumes comprise at least about 70% of perfume ingredients with relatively high boiling points (i.e. at least about 250°C) and hydrophobicities (i.e. a ClogP of at least about 3.0). Such perfume compositions may however exhibit odour characteristics which are influenced by the odour properties of the substantial quantities of enduring perfume ingredients in the perfume compositions.

EP 0003172, EP 0005618, US 4304679, US 4322308, US 4278658, US 4134838, US 4288341, US 4289641 and US 4906454 all describe perfume compositions which exhibit a deodorant action, either when applied to human skin using a cosmetically acceptable vehicle or when included in a detergent product or fabric conditioning product used in laundering of

textiles. EP 0147191 and US 4663068 describe deodorant perfume compositions which are stable in the presence of bleaching materials. A difficulty with these perfume compositions is that they generally include appreciable quantities of relatively high molecular weight perfume components which help to extend the effective lifetime of deodorant action following product application, but which tend to have less perfume impact and to exhibit odour characteristics which span a finite range.

The above described limitations of the perfume compositions of the prior art represent a compromise between long term deodorant efficacy and optimal hedonic performance. This compromise is particularly significant for perfumes characterised by "light notes" i.e. odour notes which are delicate, are not cloying and may in fact be fleeting, and dominated by what are known within the perfume industry as top-notes. By this is meant the odour notes which are detected during the first stages of perfume evaporation from a surface such as skin, and which typically, but not necessarily, tend to arise from volatile materials. Volatile materials will by their very nature evaporate faster than other perfume components and are not expected to contribute significantly to effective long term deodorant activity following application to surfaces such as skin or fabric.

The present invention concerns novel perfume compositions, particularly perfume compositions which may be described as light i.e. of delicate odour and non-enduring, yet demonstrate effective deodorant longevity.

Summary of the Invention

In one aspect, the present invention provides a perfume composition comprising:

- a) nitrile perfumery material of general formula RCN where R is a hydrocarbyl radical of formula C_xH_y , x is an integer in the range 9 to 12, and y is an integer in the range 15 to 25; and
 - b) hydrocarbon perfumery material having a boiling point of less than about $185^\circ C$ and an octanol-water partition coefficient of at least 4.0 (in logarithmic form),
- wherein a) and b) together comprise at least 20% by weight of the total weight of the composition.

Nitriles are known perfumery materials for use in deodorant perfume compositions.

For example, EP 0299561 describes bleach compositions comprising peracid-stable perfume compositions. Composition B, Composition C and Composition D of the Examples respectively describe perfume compositions comprising 0.5% of dodecyl nitrile, 0.5% of 3-methyldodecanonitrile (FRESCILE™) and 0.5% of 2-methyldecanonitrile (FRUTONILE™).

WO 00/01361 describes deodorant perfume compositions prepared from at least three of six specified categories of perfume materials; hydroxylic materials, ketones, aldehydes, ethers, esters and nitriles.

EP 1113105 relates to malodour maskant compositions containing from about 10 to about 100% by weight of 3,7-dimethyl-2-6-octadien-1-nitrile (also known as CITRALVA™) and optional components, β -naphthyl methyl ether, β -naphthyl methyl ketone, benzyl acetone, CYCLAPROP™, gamma-methyl ionone, AMBERIFF™, the ethylene glycol cyclic ester of n-docecanedioic acid, 1-cyclohexadecen-6-one and/or 1-cycloheptadecene-10-one and corn mint oil.

However, it is well known to those skilled in the art of perfumery that it can be difficult to use nitriles, particularly unsaturated nitriles such as citronellyl nitrile and geranyl nitrile, at significant levels in a perfume as they tend to smother the odour perception of other ingredients. A difficulty with some of the perfume compositions disclosed in the prior art is that they may include such components which frequently give them strong, powerful odours which are difficult for the perfumer to blend out and which can limit the usefulness of the compositions when used to perfume some other product such as a detergent composition, fabric conditioner or personal care product. As discussed above, there is generally a necessity for compromise between deodorant efficacy and acceptability as a fragrance.

The nitrile perfumery material and hydrocarbon perfumery material of the perfume compositions defined herein may be categorised as top-notes, having generally low to moderate boiling points. The materials may also be described as light notes as they

contribute delicate notes to the perfume composition which are not cloying and may in fact be fleeting.

It has been found by the inventors that the above defined perfume compositions characterised by the presence of an appreciable amount of top-note materials demonstrate unexpected high deodorant activity several hours after application. Despite the comparatively high proportion of top-note materials present, which are expected to have a relatively transient influence on body malodour perception, perfume compositions as defined herein exhibit significant deodorant activity several hours, typically at least 5 hours, after application to the body i.e. skin or to fabric. It has thus been found that perfume compositions in accordance with the present invention make it possible to obtain good deodorant performance and widely acceptable fragrance.

The term 'perfumery material' is herein taken to represent materials that may be acceptably employed within fragrances to provide an odour contribution to the overall hedonic performance of the fragrance. Typically, such materials will be generally recognised as possessing odours in their own right, and will be relatively volatile, and characterised by molecular weights within the range of around 100 to 300 amu.

The concentration of perfumery materials referred to herein is relative to the total concentration of perfumery materials present in the composition and excludes, for example, the presence of any optional solvents or diluents etc.

The octanol-water partition coefficient (P) of a material i.e. the ratio of a material's equilibrium concentration in octanol and water, is well known in the literature as a measure of hydrophobicity and water solubility (see Hansch and Leo, Chemical Reviews, 526 to 616, (1971), 71; Hansch, Quinlan and Lawrence, J.Organic Chemistry, 347 to 350 (1968), 33). High partition coefficient values are more conveniently given in the form of their logarithm to the base 10, log P. While log P values can be measured experimentally i.e. directly, and measured log P data is available for many perfumes, log P values are most conveniently calculated or approximately estimated using mathematical algorithms. There are several recognised calculation or estimation methods available commercially and/or described in the

literature (see for example A Leo, Chem.Rev 93(4), 1281-1306, (1993), "Calculating log P oct from structures"). Generally these models correlate highly but may for specific materials produce log P values which differ in absolute terms (by up to 0.5 log units or even more). However, no one model is universally accepted as the most accurate across all compounds. This is particularly true for estimates on materials of high log P (say 4 or greater). In the present specification, log P values are obtained using the estimation software commercially available as 'LogP' from Toronto-based Advanced Chemistry Development Inc (ACD) which is well-known to the scientific community, and accepted as providing high-quality predictions of log P values. References to log P values thus mean values obtained using the ACD software.

Perfume compositions in accordance with the present invention preferably comprise at least 25%, more preferably at least 30%, and most preferably about 50% by total weight of the composition of nitrile perfumery material and hydrocarbon perfumery material (ingredients (a) and (b)) as described herein.

Typically, the weight ratio of nitrile perfumery material to hydrocarbon perfumery material is in the range 10:1 to 1:2. Particularly effective perfume compositions may be prepared when the weight ratio of nitrile perfumery material to hydrocarbon perfumery material is in the range 3:1 to 2:3.

The nitrile perfumery material suitable for use herein may be saturated or unsaturated, and conveniently, the hydrocarbon perfumery material is a terpene. Examples of suitable nitrile perfumery material of category (a) and hydrocarbon perfumery material of category (b) include the following:

Table 1

Perfumery Material	Category
2-Methyldecanenitrile	(a)
3-(2,4-dimethylcyclohex-3-enylidene)propanenitrile	(a)
3,7-Dimethyl-2,6-nonadienenitrile	(a)
Camphene	(b)

Citronellyl nitrile	(a)
p-Cymene	(b)
Dodecanenitrile	(a)
Geranyl nitrile	(a)
2-tridecenenitrile	(a)
2-undecenitrile	(a)
3-methyldodecanonitrile	(a)
Limonene	(b)
Myrcene	(b)
Pinenes *	(b)
Terpinenes **	(b)
Terpinolene	(b)

* including alpha-and beta-isomers

** including alpha-, beta-, and gamma isomers

Preferred nitrile perfumery material of category (a) includes unsaturated nitrile perfumery material where when x is 9, y is 15 or 17; when x is 10, y is 17 or 19; when x is 11, y is 19 or 21 and when x is 12, y is 21 or 23. Thus, preferably, y is an integer in the range 15 to 23. A particularly preferred unsaturated nitrile perfumery material for use herein is geranyl nitrile where x is 9 and y is 15. A preferred component of the hydrocarbon perfumery material of category (b) include pinenes such as alpha-pinene and beta-pinene, or mixtures thereof.

In a preferred perfume composition herein, geranyl nitrile comprises at least 75% by weight of nitrile perfumery material and α -pinene and β -pinene, typically in a weight ratio of 2:1 to 1:2, comprise at least 75% by weight of hydrocarbon perfumery material.

Generally, perfume compositions as defined herein are non-enduring and conveniently comprise low quantities of high boiling, and highly hydrophobic perfume materials. That is, they typically comprise less than 70%, preferably less than 60% and even more preferably less than 50% of perfumery materials having a boiling point of at least about 250°C and an octanol-water partition coefficient of at least about 3.0 (in logarithmic form to base 10).

Typically, nitrile perfumery material, whether saturated or unsaturated, having at least 12 carbon atoms, will have a boiling point greater than 250°C and an octanol-water partition coefficient greater than 3.0. Such nitrile perfumery materials may be classified as 'enduring perfumery materials', which if employed in a perfume composition as defined herein are present in amount of less than 30% by weight of the perfume composition.

In a further aspect, the present invention provides a perfume composition comprising:

- a) nitrile perfumery material of general formula RCN where R is a hydrocarbyl radical of formula C_xH_y , x is an integer in the range 9 to 12, and y is an integer in the range of 15 to 25; and
 - b) hydrocarbon perfumery material having a boiling point of less than about 185°C and an octanol-water partition coefficient of at least 4.0 (in logarithmic form),
- wherein a) and b) together comprise at least 20% by weight of the total weight of the composition and wherein:
- (i) the weight ratio of nitrile perfumery material to hydrocarbon perfumery material is in the range 10:1 and 1:2; and
 - (ii) the perfume composition comprises less than 70% of perfumery materials having a boiling point of at least about 250°C and an octanol-water partition coefficient of at least 3.0 (in logarithmic form).

In the simplest case, the perfume composition consists of nitrile perfumery material and hydrocarbon perfumery material as defined herein. The perfume composition may however optionally comprise further excipients such as one or more additional perfumery materials, which will be selected so as to be congruent with the olfactory characters of the nitrile perfumery material and hydrocarbon perfumery material and to produce a fragrance having desired odour properties to meet the criteria of consumer acceptability and desired hedonic direction.

Additional perfumery materials that can be included in the perfume composition include natural products such as extracts, essential oils, absolutes, resinoids, resins, concretes etc., but also synthetic materials such as alcohols, aldehydes, ketones, ethers, acids, esters,

acetals, ketals, etc., including saturated and unsaturated compounds, aliphatic, carbocyclic, and heterocyclic compounds:

Such perfumery materials are mentioned, for example, in S. Arctander, *Perfume and Flavour Chemicals* (Montclair, N.J., 1969), in S. Arctander, *Perfume and Flavour Materials of Natural Origin* (Elizabeth, N.J., 1960) and in "Flavour and Fragrance Materials - 1991", Allured Publishing Co. Wheaton, Ill. USA.

Examples of additional perfumery materials which may be present in the perfume compositions in accordance with the invention are: acetyl cedrene, 4-acetoxy-3-pentyltetrahydropyran, 4-acetyl-6-t-butyl-1,1-dimethylindane, available under the trademark "CELESTOLIDE", 5-acetyl-1,1,2,3,3,6-hexamethylindane, available under the trademark "PHANTOLIDE", 6-acetyl-1-isopropyl-2,3,3,5-tetramethylindane, available under the trademark "TRASEOLIDE", alpha-n-amylocinnamic aldehyde, amyl salicylate, aubepine, aubepine nitrile, aurantion, 2-t-butylcyclohexyl acetate, 2-t-butylcyclohexanol, 3-(p-t-butylphenyl)propanal, 4-t-butylcyclohexyl acetate, 4-t-butyl-3,5-dinitro-2,6-dimethylacetophenone, 4-t-butylcyclohexanol, benzoin siam resinoids, benzyl acetate, benzyl propionate, benzyl salicylate, benzyl isoamyl ether, bergamot oil, bornyl acetate, butyl salicylate, carvacrol, cedar atlas oil, cedryl methyl ether, cedryl acetate, cinnamic alcohol, cinnamyl propionate, cis-3-hexenol, cis-3-hexenyl salicylate, citronella oil, citronellol, citronellyl acetate, citronellyloxyacetaldehyde, cloveleaf oil, coumarin, 9-decen-1-ol, n-decanal, n-dodecanal, decanol, decyl acetate, dihydromyrcenol, dihyromyrcenyl formate, dihydromyrcenyl acetate, dihydroterpinyl acetate, dimethylbenzyl carbinyl acetate, dimethylbenzylcarbinol, dimethylheptanol, dimethyloctanol, dimyrcetol, diphenyl oxide, ethyl naphthyl ether, ethyl vanillin, ethylene brassylate, eugenol, florocyclene, geraniol, geranium oil, geranyl acetate, 1,1,2,4,4,7-hexamethyl-6-acetyl-1,2,3,4-tetrahydronaphthalene, available under the trademark "TONALID", 1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethylcyclopenta-2-benzopyran, available under the trade mark "GALAXOLIDE", 2-n-heptylcyclopentanone, 3a,4,5,6,7,7a-hexahydro-4,7-methano-1(3)H-inden-6-ylpropionate, available under the trademark "FLOROCYCLENE", 3a-4,5,6,7,7a-hexahydro-4,7-methano-1(3)H-inden-6-ylacetate, available under the trademark "JASMACYCLENE", 4-(4'-hydroxy-4'-methylpentyl)-3-cyclohexenecarbaldehyde, alpha-

hexylcinammic aldehyde, heliotropin, hexyl aldol, hexyl cinnamic aldehyde, hexyl salicylate, hydroxycitronellal, i-nonyl formate, 3-isocamphylcyclohexanol, 4-isopropylcyclohexanol, 4-isopropylcyclohexyl methanol, indole, ionones, irones, isoamyl salicylate, isoborneol, isobornyl acetate, isobutyl salicylate, isobutylbenzoate, isobutylphenyl acetate, isoeugenol, isolongifolanone, isomethyl ionones, isononanol, isononyl acetate, isopulegol, lavandin oil, lemongrass oil, linalool, linalyl acetate, methyl beta orcinyl carboxylate (LRG 201), 1-menthol, 2-methyl-3-(p-isopropylphenyl)propanal, 2-methyl-3-(p-t-butylphenyl)propanal, 3-methyl-2-pentyl-cyclopentanone, 3-methyl-5-phenyl-pentanol, alpha and beta methyl naphthyl ketones, methyl ionones, methyl dihydrojasmonate, methyl naphthyl ether, methyl 4-propyl phenyl ether, Mousse de chene Yugo, Musk ambrette, myrtenol, neroli oil, nonanediol-1,3-diacetate, nonanol, nonanolide-1,4, nopol acetate, 1,2,3,4,5,6,7,8-octahydro-2,3,8,8-tetramethyl-2-acetyl-naphthalene, available under the trademark "ISO-E-SUPER", octanol, Oppoponax resinoid, orange oil, p-t-amylcyclohexanone, p-t-butylmethylhydrocinnamic aldehyde, 2-phenylethanol, 2-phenylethyl acetate, 2-phenylpropanol, 3-phenylpropanol, para-menthan-7-ol, para-t-butylphenyl methyl ether, patchouli oil, pelargene, petitgrain oil, phenoxyethyl isobutyrate, phenylacetaldehyde diethyl acetal, phenylacetaldehyde dimethyl acetal, phenylethyl n-butyl ether, phenylethyl isoamyl ether, phenylethylphenyl acetate, pimento leaf oil, rose-d-oxide, Sandalone, styrallyl acetate, 1,1,4,4-tetramethyl-6-acetyl-7-ethyl-1,2,3,4-tetrahydronaphthalene, available under the trademark "VERSALIDE", 3,3,5-trimethyl hexyl acetate, 3,5,5-trimethylcyclohexanol, terpeneol, terpinyl acetate, tetrahydrogeraniol, tetrahydrolinalool, tetrahydromuguol, tetrahydromyrcenol, thyme oil, trichloromethylphenylcarbinyl acetate, tricyclodecenyl acetate, tricyclodecenyl propionate, 10-undecen-1-al, gamma undecalactone, 10-undecen-1-ol, undecanol, vanillin, vetiverol, vetiveryl acetate, vetyvert oil, acetate and propionate esters of alcohol in the list above, aromatic nitromusk fragrances, indane musk fragrances, isochroman musk fragrances, macrocyclic ketones, macrolactone musk fragrances, and tetralin musk fragrances.

Further excipients which may be included in the perfume compositions of the present invention include various solvents, diluents or vehicles for other materials which are known for perfumery use and typically have a low odour or no odour. Examples of suitable solvents include benzyl alcohol, benzyl benzoate, dipropylene glycol, triacetin, Herculyn

DTM, diethyl phthalate, triacetin, isopropyl myristate, triethyl citrate and acetyl tributyl citrate.

In an even further aspect the perfumes comprise compositions in which at least 20% by weight of the composition is constituted by perfume materials from both of the following categories:

- a) perfumery nitriles of general formula RCN wherein R is a hydrocarbyl radical of formula C_xH_y, where 'x' is an integer in the range 10 to 12, and 'y' is an integer in the range 15 to 23;
- b) perfumery hydrocarbons characterised by boiling points of less than 185°C and octanol-water partition coefficients of at least 4.0 (logarithm base 10);

provided also that:

- (i) the ratio of class a) ingredients to class b) ingredients lies within the range 10:1 to 1:2 (by weight);
- (ii) the composition comprises less than 70% of materials that have boiling points of at least 250°C and octanol-water partition coefficients of at least 3.0 (logarithm base 10).

The compositions of the present invention can be used in or constitute a wide range of products. The invention in a further aspect therefore provides a product comprising a perfume composition in accordance with the invention.

Suitable products include, but are not limited to, products which provide a deodorant action when applied to the body e.g. deodorants and antiperspirants including different physical forms such as roll ons, gels, sticks, and aerosols; other personal products such as deocolognes, talcum powders, hand creams, lotions, skin and hair conditioners, sunscreens, soaps, shampoos, and shower gels; and laundry and household products such as laundry powders, laundry liquids, laundry tablets, rinse conditioners and fabric treatment products including fabric refresher products e.g. sprays, starch sprays, ironing sprays and stain remover sprays.

A preferred product is a laundry product and more particularly a laundry powder. Thus, in a preferred embodiment herein, the laundry product, particularly laundry powder, comprises between 0.01% to 10% and more preferably 0.1% to 1.0% by weight of the laundry product of a perfume composition in accordance with the present invention.

Perfume compositions in accordance with the present invention, in their simplest form as a fragrance, or as part of a product, may be applied to a surface including skin, hair, fabric etc, directly or indirectly. For example, when a perfume composition is part of a laundry product for washing fabric the composition may be applied to the fabric indirectly, i.e. via the wash liquor during the wash process.

The physical properties of the perfume materials of categories (a) and (b) are such that they are likely to rapidly evaporate from the surface they are applied to. The actual rate of evaporation is dependent upon a number of factors including the environmental conditions prevalent during application and following application e.g. temperature of the surface, the nature of the surface, and the presence and nature of other materials which may be delivered together with the fragrance during product treatment e.g. surfactant actives. Without wishing to be bound or limited by theory, it is thought that odour synergies between the nitrile perfumery material and hydrocarbon perfumery material may play a part in delivering efficient body malodour counteraction even at low residual levels.

Perfume compositions of the invention may also be incorporated into textiles directly during manufacture using techniques known in the art, to provide long lasting deodorant protection. It is also known in the art to carry or to encapsulate perfumes within other materials such as porous solids or polymeric matrices, in order to provide extended lifetimes, and to provide the possibility of triggered release, for example, during perspiration. Such techniques are applicable within the scope of the present invention.

In a further aspect, the invention provides a method of counteracting body malodour on a surface, by applying to the surface, a perfume composition in accordance with the present invention.

A perfume composition in accordance with the present invention counteracts body malodour on a surface, typically by reducing the perceived intensity of the body malodour evolved from the surface, or inhibiting or suppressing the olfactory detection of the evolution of body malodour from the surface when it comes into contact with body malodour, as appropriate.

In an even further aspect, the invention provides the use, as a deodorant, of a perfume composition in accordance with the present invention.

The invention is illustrated by the following examples.

All percentages are by weight unless otherwise indicated.

Example 1

The following perfume, Perfume A, was prepared in accordance with the present invention and in the proportions indicated in Table 2 below:

Table 2 - Perfume A

Ingredient	w/w%	Category
alpha-Pinene	5	b
alpha-Terpineol	9.5	-
Benzyl benzoate	5.0	-
beta-Pinene	5.0	b
Camphene	0.8	b
Cineole	1.0	-
Dihydromyrcenol	25.0	-
Dodecanal (50% in DPG)	1.0	-
Geranyl nitrile	15.0	a
Lemongrass oil	3.0	-
Lilial	0.5	-
Linalol	6.1	-
Linalyl acetate	9.2	-

Litsea cubeba	7.2	-
Methyl chavicol	1.0	-
Octanal (50% in DPG)	1.5	-
Ortho-tert.-butylcyclohexanol	0.7	-
para-Cymene	0.9	b
para-tert-butyl cyclohexyl acetate	0.3	-

Sum class (a) = 15.0% Sum class (b) = 11.7%

Perfume A was tested for deodorant action in a laundry powder using the Malodour Reduction Value test as generally described in EP 545556. The results of this 5 hour deodorant test are given in Table 3, and demonstrate that Perfume A surprisingly provides significant deodorant action many hours after its application to cloth.

Table 3 Results of 5 hr Malodour Reduction deodorant efficacy test

Average panel score Perfume A	2.28
Control panel score	3.23
Malodour Reduction Value Perfume A	0.95
Malodour Reduction Value as percentage of control score	29.4%
Difference for significance @ 99% = 0.23.	

Example 2

The Malodour Reduction Value test discussed above was repeated for Perfume A described in Example 1 above. Perfume A in this test was labelled Perfume 4500A.

Comparative testing was also carried out under the same conditions and using the same protocol with perfumes based on Perfume A, but modified to fall outside the scope of the invention. The comparative perfumes tested were as follows:

Perfume 4500B containing 50% of the amount of geranyl nitrile of Perfume A;

Perfume 4500D containing no nitrile perfumery material; and

Perfume 4500C containing 25% of the amount of hydrocarbon perfumery materials (category (b) materials) of Perfume A.

The results of the 5 hour deodorant test are given in Table 4.

Table 4 Results of 5 hr Malodour Reduction deodorant efficacy test

Average panel score Perfume 4500A	2.48
Average panel score Perfume 4500B	2.57
Average panel score Perfume 4500D	2.59
Average panel score Perfume 4500C	2.65

It will be noted that the average panel score for Perfume 4500A is higher than for the test described in Example 1. This is likely to arise because of noise changes during test conditions (e.g. humidity, temperature etc.) which may affect deodorancy (from both the perspectives of malodour generation and malodour counteraction). Nevertheless, the results clearly show that Perfume 4500A (Perfume A) in accordance with the present invention demonstrates good deodorant action and that the modifications made to the formulation adversely affected deodorant action and efficacy.

Claims

1. A perfume composition comprising:
 - a) nitrile perfumery material of general formula RCN where R is a hydrocarbyl radical of formula C_xH_y , x is an integer in the range 9 to 12, and y is an integer in the range 15 to 25; and
 - b) hydrocarbon perfumery material having a boiling point of less than about $185^{\circ}C$ and an octanol-water partition coefficient of at least 4.0 (in logarithmic form),wherein a) and b) together comprise at least 20% by weight of the total weight of the composition.
2. A perfume composition according to claim 1, comprising at least 25%, preferably at least 30% by total weight of the perfume composition of nitrile perfumery material and hydrocarbon perfumery material.
3. A perfume composition according to claim 1 or 2, wherein the weight ratio of nitrile perfumery material to hydrocarbon perfumery material is in the range 10:1 to 1:2, preferably 3:1 to 2:3.
4. A perfume composition according to any one of the preceding claims, wherein the nitrile perfumery material is 2-methyldecanenitrile, 3-(2,4-dimethylcyclohex-3-enylidene)propanenitrile, 3,7-dimethyl-2,6-nonadienenitrile, citronellyl nitrile, 2-tridecenenitrile, 2-undecenitrile, 3-methyldodecanonitrile, dodecanenitrile or geranyl nitrile, and mixtures thereof.
5. A perfume composition according to any one of the preceding claims, wherein the hydrocarbon perfumery material is a terpene.
6. A perfume composition according to any one of the preceding claims, wherein the hydrocarbon perfumery material is camphene, p-cymene, limonene, myrcene, α -pinene, β -pinene, α -terpinene, β -terpinene, γ -terpinene or terpinolene, and mixtures thereof.

7. A perfume composition according to any one of the preceding claims, wherein the perfume composition comprises less than 70% of perfumery materials having a boiling point of at least about 250°C and an octanol-water partition coefficient of at least 3.0 (in logarithmic form).

8. A perfume composition according to any one of the preceding claims, wherein the nitrile perfumery material is unsaturated, and x is an integer in the range 9 to 12, and y is an integer in the range 15 to 23.

9. A perfume composition according to any one of the preceding claims, wherein x is 9 and y is 15, and the nitrile perfumery material is geranyl nitrile.

10. A perfume composition according to any one of the preceding claims, wherein geranyl nitrile comprises at least 75% by weight of nitrile perfumery material.

11. A perfume composition according to any one of the preceding claims, wherein the hydrocarbon perfumery material comprises a mixture of α -pinene and β -pinene.

12. A perfume composition according to claim 11, wherein α -pinene and β -pinene comprise at least 75% by weight of hydrocarbon perfumery material.

13. A product comprising a perfume composition in accordance with any one of claims 1 to 12.

14. A laundry product comprising between 0.01% to 10% by weight of the laundry product of a perfume composition in accordance with any one of claims 1 to 12.

15. A perfume composition in which at least 20% by weight of the composition is constituted by perfume materials from both of the following categories:

a) perfumery nitriles of general formula RCN wherein R is a hydrocarbyl radical of formula C_xH_y, where 'x' is an integer in the range 10 to 12, and 'y' is an integer in the range 15 to 23;

b) perfumery hydrocarbons characterised by boiling points of less than 185°C and octanol-water partition coefficients of at least 3.0 (logarithm, base 10);

provided also that:

- (i) the ratio of class a) ingredients to class b) ingredients lies within the range 10:1 to 1:2 (by weight);
- (ii) the composition comprises less than 70% of materials that have boiling points of at least 250°C and octanol-water partition coefficients of at least 3.0 (logarithm base 10).

Inter. onal Application No

A. CLASSIFICATION OF SUBJECT MATTER

According to International Patent Classification (IPC) or to both national classification and IPC

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

^o Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

18 July 2002

24/07/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer _____

Rasmusson, R

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/01609

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 6 086 903 A (TRINH ET AL) 11 July 2000 (2000-07-11) column 3, line 1-6 column 5, line 1 - line 49 table 2 example I	1-15
A	WO 93 04158 A (PROCTER & GAMBLE) 4 March 1993 (1993-03-04) claims 1,5,6,10	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/GB 02/01609

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6024943	A	15-02-2000	AT 213001 T	15-02-2002
			AU 729041 B2	25-01-2001
			AU 5331798 A	17-07-1998
			AU 730956 B2	22-03-2001
			AU 5331898 A	17-07-1998
			AU 5331998 A	17-07-1998
			BR 9713619 A	11-04-2000
			BR 9713634 A	11-04-2000
			DE 69710367 D1	21-03-2002
			DK 950070 T3	13-05-2002
			EP 0950087 A1	20-10-1999
			EP 0950088 A1	20-10-1999
			EP 0950070 A1	20-10-1999
			WO 9828396 A1	02-07-1998
			WO 9828398 A1	02-07-1998
			WO 9828339 A1	02-07-1998
			JP 2001507058 T	29-05-2001
			JP 2001507384 T	05-06-2001
			JP 2001507059 T	29-05-2001
			US 6194375 B1	27-02-2001
			US 6329057 B1	11-12-2001
			ZA 9711578 A	23-06-1999
			ZA 9711582 A	23-06-1999
			ZA 9711589 A	23-06-1999
WO 9900025	A	07-01-1999	AU 8169798 A	19-01-1999
			AU 8169898 A	19-01-1999
			BR 9810339 A	05-09-2000
			BR 9810935 A	26-09-2000
			CN 1268873 T	04-10-2000
			CN 1268874 T	04-10-2000
			EP 0991327 A1	12-04-2000
			EP 0991328 A1	12-04-2000
			HU 0003605 A2	28-05-2001
			HU 0004533 A2	28-04-2001
			JP 2002507236 T	05-03-2002
			JP 2002511758 T	16-04-2002
			TR 9903224 T2	21-07-2000
			TR 9903231 T2	21-07-2000
			WO 9900025 A1	07-01-1999
			WO 9900026 A1	07-01-1999
			ZA 9805560 A	28-12-1998
			ZA 9809648 A	23-04-1999
US 6001789	A	14-12-1999	CA 2249293 A1	25-09-1997
			EP 0888446 A1	07-01-1999
			JP 2000502744 T	07-03-2000
			WO 9734993 A1	25-09-1997
US 6086903	A	11-07-2000	BR 9708304 A	03-08-1999
			CA 2246293 A1	28-08-1997
			EP 0886516 A1	30-12-1998
			WO 9730689 A1	28-08-1997
WO 9304158	A	04-03-1993	AT 155164 T	15-07-1997
			AU 2473592 A	16-03-1993
			CA 2115539 A1	04-03-1993
			CN 1071451 A	28-04-1993

INTERNATIONAL SEARCH REPORT

Information on patent family members.

Inter. Application No

PCT/GB 02/01609

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9304158	A	DE 69220803 D1	14-08-1997
		DE 69220803 T2	18-12-1997
		EP 0601020 A1	15-06-1994
		ES 2103960 T3	01-10-1997
		JP 6510077 T	10-11-1994
		MX 9204867 A1	01-04-1993
		NZ 244032 A	27-02-1996
		PT 100806 A	30-09-1993
		TR 27092 A	18-10-1994
		WO 9304158 A1	04-03-1993
		US 5614484 A	25-03-1997